

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

98889-CN

ID	Weight %	Concentration (mg/mL)	
D9-THC	0.218	2.03	
THCV	ND	ND	
CBD	3.76	35.0	
CBDV	<loq< td=""><td><loq< td=""><td></td></loq<></td></loq<>	<loq< td=""><td></td></loq<>	
CBG	0.0731	0.681	
CBC	0.183	1.70	
CBN	<loq< td=""><td><loq< td=""><td></td></loq<></td></loq<>	<loq< td=""><td></td></loq<>	
THCA	0.0564	0.525	
CBDA	3.62	33.7	
CBGA	0.0766	0.713	
D8-THC	ND	ND	
exo-THC	ND	ND	
Total	8.00	74.5	0% Cannabinoids (wt%) 3.76%
Max THC	0.268	2.49	Limit of Quantitation (LOQ) = 0.0113 wt%
Max CBD	6.93	64.5	Limit of Detection (LOD) = 0.0038 wt%

Ratio of Total CBD to THC 25.9:1

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: MAX THC = $(0.877 \times THCA) + THC$. This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND=None detected above the limits of detection (LOD), which is one third of Limit of Quantification (LOQ). For values reported as "<LOQ", the estimated value is included in the calculated Total.

Test Date: 11/1/2021

TP: Terpenes Profile [WI-10-27]

Client sample analysis was performed using full evaporative technique (FET) headspace sample delivery and gas chromatographic (GC) compound separation. A combination of flame ionization detection (FID) and/or mass spectrometric (MS) detection with mass spectral confirmation against the National Institute of Standards and Technology (NIST) Mass Spectral Database, Revision 2017 were used. Chromatographic and/or mass spectral data were processed by quantitatively comparing the analytical peak areas against calibration curves prepared from certified reference standards.

Analyst: CJS

98889-TP

Compound	CAS	Conc. (wt%)	Conc. (ppm)) Qualitative Profile
alpha-pinene	80-56-8	0.0171	171	
camphene	79-92-5	0.0006	5.66	
sabinene*	3387-41-5	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
beta-myrcene	123-35-3	0.0122	122	
beta-pinene	127-91-3	0.0043	43.0	
alpha-phellandrene	99-83-2	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
alpha-terpinene	99-86-5	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
D-limonene	138-86-3	0.0032	32.4	
p-cymene	99-87-6	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
eucalyptol	470-82-6	0.0054	53.8	
gamma-terpinene	99-85-4	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
terpinolene	586-62-9	<rl< td=""><td><rl< td=""><td></td></rl<></td></rl<>	<rl< td=""><td></td></rl<>	
linalool	78-70-6	0.0095	95.2	
L-fenchone*	7787-20-4	0.0018	18.4	
beta-caryophyllene	87-44-5	0.0465	465	
alpha-humulene	6753-98-6	0.0088	88.0	
trans-nerolidol	40716-66-3	0.0014	14.3	
guaiol	489-86-1	0.0097	97.4	
caryophyllene oxide	1139-30-6	0.0018	17.7	
alpha-bisabolol	23089-26-1	0.0118	118	
			ppm 0	0.00 250.00 500.0

Total Terpene: 0.1 wt%

* Certified reference standard not available for this compound. Concentration is estimated using the response factor from alpha-pinene. ND = None Detected. RL = Reporting Limit of 5 ppm.

END OF REPORT