

# Mepilex® Border Lite



# THE SELF-ADHERENT THIN BORDERED FOAM DRESSING WITH SAFETAC® TECHNOLOGY

- Minimizes pain and trauma at dressing changes<sup>1,2,3</sup>
- Excellent and effective exudate management

#### Safetac® technology layer

- Reduces pain and trauma during dressing change<sup>1</sup>
- Does not adhere to the moist wound bed but to dry skin only<sup>3</sup>
- Seals the wound margins and reduces risk of maceration4

#### 2-layer pad

- Absorbs exudate
- Maintains integrity

#### Polyurethane backing film


- High MVTR
- Moisture proof
- Bacteria and viral barrier<sup>†</sup>



#### Safetac® technology

### Safetac® technology. Less pain and less trauma.


Safetac® is a patented adhesive technology that minimizes pain to patients and trauma to wounds. Safetac® technology is available exclusively on Mölnlycke Health Care dressings, including Mepilex® wound care dressings, Mepitel®, Mepiform® and Mepitac®.





No skin stripping with Safetac® technology

## How to use Mepilex® Border Lite



Clean the wound area. Dry the surrounding skin thoroughly. Skin barrier products not necessary. Remove the first release film.



Apply the adherent side to the wound. Do not stretch. Remove the second release film. Smooth down border



### How Mepilex® Border Lite works

Mepilex® Border Lite is a thin and conformable version of Mepilex® Border, ideal for body contours and hard to dress areas and is available in smaller sizes. It is a self-adherent dressing that absorbs exudate, maintains a moist wound environment and minimizes the risk of peri-wound maceration. The Safetac® layer allows the dressing to be changed without causing additional pain to the patient or trauma to the wound and surrounding skin.

#### Benefits of Mepilex® Border Lite


- Minimizes trauma to the wound and pain to the patient during dressing changes
- Excellent fit to the wound and body contours
- A thin and conformable all-in-one dressing
- Broad selection of sizes
- Barrier to outside moisture, viruses and bacteria
- · Minimizes the risk of maceration
- · Does not adhere to moist wound
- Adheres gently and securely to dry, intact surrounding skin

Wear time: Up to 7 days

Note: In case of clinical signs of infection the use of Mepilex® Border Lite may be continued if proper infection treatment is initiated.

#### Areas of use

- Non to low exuding wounds
- Leg and foot ulcers
- Pressure ulcers
- Traumatic wounds (e.g., abrasions, cuts, finger injuries, blisters, and skin tears)
- Protection of compromised and/or fragile skin





Perfect for hard to dress areas and body contours.

Mepilex® Border Lite Ordering Information‡				
Product Code	Size	Pieces/ box	Pieces/ case	HCPCS
281000	1.6" x 2" (4 x 5 cm)	10	70	_
281100	2" x 5" (5 x 12.5 cm)	5	65	A6212
281200	3" x 3" (7.5 x 7.5 cm)	5	70	A6212
281300	4" x 4" (10 x 10 cm)	5	50	A6212
281500	6" x 6" (15 x 15 cm)	5	50	A6213

#### References

- 1. White R., A Multinational survey of the assessment of pain when removing dressings. Wounds UK 2008; Vol 4, No 1.
- 2. Dykes PJ et al. Effects of adhesive dressings on the stratum corneum of the skin. J Wound Care 2001; 10(2):7-10.
- Dykes PJ. The effect of adhesive dressing edges on cutaneous irritancy and skin barrier function.
  J Wound Care 2007. 16(3):97-100.
- Meaume S et al. A study to compare a new self adherent soft silicone dressing with a self adherent polymer Management, 2003.
- 5. External lab report: NAMSA 09C 29253 01/09C 29253 02.
- † Microbes larger than 25 nm
- ‡ Packaged sterile in single packs



